

A NOTE ABOUT RATIONAL REPRESENTATIONS OF DIFFERENTIAL GALOIS GROUPS.

MARC REVERSAT

ABSTRACT. We give a description of the rational representations of the differential Galois group of a Picard-Vessiot extension.

CONTENTS

1. On representations of Galois groups of Picard-Vessiot extensions.	1
2. A correspondance.	6
References	8

We give a description of the rational representations of the differential Galois group of a Picard-Vessiot extension (theorems 1.1 and 2.1). This gives a new description of the differential Galois correspondence. More results will be given for abelian differential extensions in a forthcoming paper [1], especially the analog of the Artin correspondence.

1. ON REPRESENTATIONS OF GALOIS GROUPS OF PICARD-VESSIOT EXTENSIONS.

In all this note we consider differential fields with algebraically closed fields of constants, denoted by C . The derivative will be denoted by a dash.

Let K be a differential field. Let $n \geq 1$ be an integer, $M_n(K)$ and $GL_n(K)$ are the usual notations for algebra and group of $n \times n$ matrices with entries in K . The group $GL_n(K)$ acts on $M_n(K)$ by the following rule:

$$\begin{aligned} GL_n(K) \times M_n(K) &\longrightarrow M_n(K) \\ (U, A) &\longmapsto U'U^{-1} + UAU^{-1} \end{aligned}$$

where if $U = (u_{i,j})_{1 \leq i,j \leq n}$, then $U' = (u'_{i,j})_{1 \leq i,j \leq n}$. This action can be defined in an other way, maybe more comprehensible. Consider the group

$$H_n(K) := M_n(K) \times GL_n(K),$$

Date: February 2, 2008.

the law being defined by the following formula: for all A, B in $M_n(K)$ and all $F, G \in GL_n(K)$

$$(A, F)(B, G) = (A + FBF^{-1}, FG).$$

It admits the subgroups

$$\Delta_n(K) := \{(U'U^{-1}, U) \mid U \in GL_n(K)\},$$

$\{0\} \times GL_n(K)$ and $M_n(K) \times \{1\}$, this last one being normal. We set

$$Z_n(K) := \Delta_n(K) \backslash H_n(K) / (\{0\} \times GL_n(K)).$$

The group $GL_n(K)$ acts on $M_n(K) \times \{1\}$ by the following rule:

$$\begin{aligned} GL_n(K) \times (M_n(K) \times \{1\}) &\longrightarrow (M_n(K) \times \{1\}) \\ (U, (A, 1)) &\longmapsto (U'U^{-1}, U)(A, 1)(0, U^{-1}) \\ &= (U'U^{-1} + UAU^{-1}, 1) \end{aligned}$$

With the identification $M_n(K) = (M_n(K) \times \{1\})$ and inclusion $(M_n(K) \times \{1\}) \subset H_n(K)$, it induces a canonical bijection

$$(1) \quad GL_n(K) \backslash M_n(K) \simeq \Delta_n(K) \backslash H_n(K) / (\{0\} \times GL_n(K)) = Z_n(K).$$

We will use below these two definitions of $Z_n(K)$.

Let A be in $M_n(K)$ and F be in $GL_n(K)$, we denote by $[(A, F)]$, resp. $[A]$, the class in $Z_n(K)$ of $(A, F) \in H_n(K)$, resp. of $A \in M_n(K)$.

Let L/K be a Picard-Vessiot extension. The inclusion $H_n(K) \subseteq H_n(L)$ gives rise to a map $\alpha(L/K) : Z_n(K) \rightarrow Z_n(L)$. We set

$$Z_n(L/K) := \{a \in Z_n(K) \mid \alpha(L/K)(a) = [0]\}.$$

For any group G we denote by $\text{Rep}_n(G)$ the set of equivalent classes of representations of G in $GL_n(C)$, if $G = d\text{Gal}(L/K)$ is the differential group of L over K , we set $\text{Rep}_n(L/K) := \text{Rep}_n(G)$.

Theorem 1.1. *Let L/K be a Picard-Vessiot extension, then there exists a natural bijection between $Z_n(L/K)$ and $\text{Rep}_n(L/K)$.*

Proof. First of all we recall some facts that are of main importance in our proof.

1.0.1. *The representation c_A .* Consider the differential equation $Y' = AY$, with $A \in M_m(K)$, and let E/K be a corresponding Picard-Vessiot extension, i.e. E is generated over K by the coefficients of a fundamental matrix F_A of the equation. The rational representation c_A is

$$\begin{aligned} d\text{Gal}(E/K) &\longrightarrow GL_m(C) \\ \sigma &\longmapsto c_A(\sigma) \end{aligned}$$

where $c_A(\sigma)$ is such that $\sigma(F_A) = F_A c_A(\sigma)$. Note that c_A depends only on the class $[A]$ of $(A, 1)$ in $Z_m(K)$, because if $B = U'U^{-1} + UAU^{-1}$ with $U \in GL_m(K)$, a fundamental matrix of the equation $Y' = BY$ is UF_A and we see that $c_B = c_A$. Note also that an other fundamemtal matrix is of the form $F_A \gamma$, with $\gamma \in GL_m(C)$, then it gives the representation $\gamma^{-1} c_A \gamma$ equivalent to c_A .

We will write equivalently c_A , $c_{[A]}$ or c_E for this class of representations.

1.0.2. *The Galois group $dGal(E/K)$.* Let $A \in M_m(K)$ and

$$R = K[(X_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] / q = K[(x_{i,j})_{1 \leq i,j \leq m}]$$

be a Picard-Vessiot ring over K for the equation $Y' = AY$. In these formulas the $X_{i,j}$ are indeterminates, the ring $K[(X_{i,j})_{1 \leq i,j \leq m}]$ is equipped by the derivation satisfying $(X'_{i,j})_{1 \leq i,j \leq m} = A(X_{i,j})_{1 \leq i,j \leq m}$, “det” is the determinant of the matrix $(X_{i,j})_{1 \leq i,j \leq m}$, q is a maximal differential ideal and $x_{i,j}$ is the image of $X_{i,j}$. Let $E = \text{Quot}(R)$ and $\mathfrak{U} = dGal(E/K)$, consider

$$\begin{aligned} K[(X_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] &\subseteq E[(X_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] \\ &= E[(Y_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] \supseteq C[(Y_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}], \end{aligned}$$

where $(Y_{i,j})_{1 \leq i,j \leq m}$ is defined by $(X_{i,j})_{1 \leq i,j \leq m} = (x_{i,j})_{1 \leq i,j \leq m} (Y_{i,j})_{1 \leq i,j \leq m}$. Note that $Y'_{i,j} = 0$. We know that

$$\mathfrak{U} = dGal(E/K) = \text{Spec} C[(Y_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] / J$$

where $J = qE[(Y_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] \cap C[(Y_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}]$ ([2] proof of prop. 1.24 or the beginning of §1.5). We denote by $y_{i,j}$ the image of $Y_{i,j}$, then we have

$$\mathfrak{U} = dGal(E/K) = \text{Spec} C[(y_{i,j})_{1 \leq i,j \leq m}]$$

1.0.3. $\mathfrak{U} = dGal(E/K)$ as a torsor. We continue with the previous notations. Set $\mathcal{T} = \text{Spec} R$, we know that $\mathcal{T}$ is an $\mathfrak{U}$ -torsor over K ([2] theorem 1.30), moreover, we know that there exists a finite extension $\tilde{K}$ of K such that $\mathcal{T} \times_K \tilde{K} = \text{Spec}(R \otimes_K \tilde{K})$ is a trivial $\mathfrak{U}$ -torsor over $\tilde{K}$ ([2] cor. 1.31), this means that there exists $\underline{b} \in \mathcal{T}(\tilde{K})$ such that the following map is an isomorphism of $\tilde{K}$ -schemes

$$\begin{aligned} \psi : \quad \mathfrak{U} \times_C \tilde{K} &\longrightarrow \mathcal{T} \times_K \tilde{K} \\ (c_{i,j})_{1 \leq i,j \leq m} &\longmapsto \underline{b}(c_{i,j})_{1 \leq i,j \leq m} \end{aligned}$$

($\underline{b}$ can be seen as a matrix, on the right this is a product of matrices; see the definition of R above).

1.0.4. *Galois actions.* Let σ be an element of $\mathfrak{U} = dGal(E/K)$, the action of σ on R is given by the images of the $x_{i,j}$, which are defined by the matrix formula $(\sigma(x_{i,j})) = (x_{i,j})_{c_E(\sigma)}$. We denote by σ^b the morphism induces by σ on $\mathcal{T}$ or on $\mathcal{T} \times_K \tilde{K}$, this is the action of $\mathfrak{U}$ which defines the torsor structure. An element of $\mathcal{T}(\tilde{K})$ can be represented by a matrix $\underline{a} = (a_{i,j})_{1 \leq i,j \leq m}$ with $a_{i,j}$ in $\tilde{K}$, its image is $\sigma^b(\underline{a}) = \underline{a}_{c_E(\sigma)}$. For any σ in $\mathfrak{U}$ denote by λ_σ the right translation on $\mathfrak{U}$ by σ , i.e.

$$\begin{aligned} \lambda_\sigma : \quad \mathfrak{U} &\longrightarrow \mathfrak{U} \\ \tau &\longmapsto \tau\sigma \end{aligned}$$

Write again λ_σ for $\lambda_\sigma \times \text{Id}_{\tilde{K}} : \mathfrak{U} \times_C \tilde{K} \rightarrow \mathfrak{U} \times_C \tilde{K}$, then the morphism ψ of (1.0.3) is equivariant, this means that for any $\sigma \in \mathfrak{U}$ the following diagram is commutative

$$\begin{array}{ccc} \mathfrak{U} \times_C \tilde{K} & \xrightarrow{\psi} & \mathfrak{T} \times_K \tilde{K} \\ \lambda_\sigma \downarrow & & \downarrow \sigma^\flat \\ \mathfrak{U} \times_C \tilde{K} & \xrightarrow{\psi} & \mathfrak{T} \times_K \tilde{K} \end{array}$$

Proof of the theorem, the map $Z_n(L/K) \rightarrow R_n(L/K)$.

Let $A \in M_n(K)$ such that $[A] \in Z_n(L/K)$, then there exists $U \in \text{GL}_n(L)$ such that

$$A = U'U^{-1},$$

this means that U is a fundamental matrix of the equation $Y' = AY$, as it is with entries in L , it exists a differential subextension E of L which is a Picard-Vessiot extension for the equation $Y' = AY$. Denote by ρ_A the representation

$$(2) \quad \rho_A : d\text{Gal}(L/K) \xrightarrow{\text{restriction}} d\text{Gal}(E/K) \xrightarrow{c_A} \text{GL}_n(C).$$

Now we prove that this representation ρ_A does not depend on the class of A in $Z_n(K)$ and of the choice of $U \in \text{GL}_n(L)$ such that $A = U'U^{-1}$.

Let $B \in M_n(K)$ such that $[B] = [A]$ in $Z_n(K)$, then there exists $W, T \in \text{GL}_n(K)$ such that

$$(B, 1) = (W'W^{-1}, W)(A, 1)(0, T),$$

it follows that $B = W'W^{-1} + WAW^{-1}$, this means that WU is a fundamental matrix of the equation $Y' = BY$. We see that $\rho_A = \rho_B$, and we denote this representation by $\rho_{[A]}$.

Let $V \in \text{GL}_n(L)$ such that $A = U'U^{-1} = V'V^{-1}$, then we see that $(V^{-1}U)' = 0$, this means that there exists $\gamma \in \text{GL}_n()$ such that $U = V\gamma$ and the two representations define as before in (2) are conjugate.

Then to each element $[A]$ of $Z_n(L/K)$ we have associated the element $\rho_{[A]}$ of $\text{Rep}_n(L/K)$.

Proof of the theorem, the map $R_n(L/K) \rightarrow Z_n(L/K)$.

Let $\rho : d\text{Gal}(L/K) \rightarrow \text{GL}_n(C)$ be a rational representation. Let E be the fixed field of $\ker \rho$, we set $\mathfrak{U} = d\text{Gal}(E/K)$ and we denote again by ρ the representation $\mathfrak{U} \hookrightarrow \text{GL}_n(C)$ coming from the given one. The field E is a Picard-Vessiot extension corresponding to an equation $Y' = AY$, with $A \in M_n(K)$. Our aim is to prove that one can chose A in $M_n(K)$, i.e. $m = n$, and that this gives the inverse map of $[A] \mapsto \rho_{[A]}$.

We use the previous notations and descriptions of E , R , $\mathfrak{U}$, $\mathfrak{T}$ etc. We set $\text{GL}_n(C) = \text{Spec}C[(T_{r,s})_{1 \leq r,s \leq n}, (\det)^{-1}]$, let

$$\rho^\sharp : C[(T_{r,s})_{1 \leq r,s \leq n}, (\det)^{-1}] \longrightarrow C[(Y_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] / J$$

be the comorphism of $\rho : \mathfrak{U} \hookrightarrow \mathrm{GL}_n(C)$; $\rho^\sharp$ is onto. Set $I = \ker(\rho^\sharp)$, then we have an isomorphism induces by $\rho^\sharp$

$$\bar{\rho} : C[(T_{r,s})_{1 \leq r,s \leq n}, (\det)^{-1}] / I \simeq C[(Y_{i,j})_{1 \leq i,j \leq m}, (\det)^{-1}] / J.$$

Let $t_{r,s}$ be the image of $T_{r,s}$ in the quotient on the left, and recall that $y_{i,j}$ are that of $Y_{i,j}$ in the quotient on the right, then the preceding formula can be written

$$\bar{\rho} : C[(t_{r,s})_{1 \leq r,s \leq n}] \simeq C[(y_{i,j})_{1 \leq i,j \leq m}].$$

Set $\mathfrak{V} = \mathrm{Spec}(C[(t_{r,s})_{1 \leq r,s \leq n}])$, this is an algebraic subgroup of $\mathrm{GL}_n(C)$, it is isomorphic to $\mathfrak{U}$ via the morphism induces by $\bar{\rho}$, denoted by abuse of language $\rho : \mathfrak{U} \simeq \mathfrak{V}$.

The composed morphism (see (1.0.3))

$$(3) \quad \varphi : \mathcal{T} \otimes_K \tilde{K} \xrightarrow{\psi^{-1}} \mathfrak{U} \times_C \tilde{K} \xrightarrow{\rho \times \mathrm{Id}_{\tilde{K}}} \mathfrak{V} \times_C \tilde{K}$$

is an isomorphism of $\tilde{K}$ -schemes, equivariant for the actions of $\mathfrak{U}$ and $\mathfrak{V}$, this means that for any σ in $\mathfrak{U}$ we have $\varphi \circ \sigma^b = \lambda_{\rho(\sigma)} \circ \varphi$, where, as before, $\lambda_{\rho(\sigma)}$ is the endomorphism of $\mathfrak{V} \times_C \tilde{K}$ coming from the right translation by $\rho(\sigma)$ on $\mathfrak{V}$ (1.0.4).

Lemma 1.2. *Let $\varphi^\sharp$ be the comorphism of φ (see (2)) and for any $r, s = 1, \dots, n$ set $z_{r,s} = \varphi^\sharp(t_{r,s})$ (recall that $\mathfrak{V} = \mathrm{Spec} C[(t_{r,s})_{1 \leq r,s \leq n}]$). Then, for all $\sigma \in \mathfrak{U}$, there exists a matrix $a(\sigma) \in \mathrm{GL}_n(C)$ such that we have the equality of matrices: $(\sigma(z_{r,s}))_{1 \leq r,s \leq n} = (z_{r,s})_{1 \leq r,s \leq n} a(\sigma)$.*

Proof. Denote by $\lambda_{\rho(\sigma)}^\sharp$ the comorphism of the right translation by $\rho(\sigma)$ on $\mathfrak{V} \times_C \tilde{K}$, we have the equalities of matrices

$$\begin{aligned} (\sigma(z_{r,s}))_{1 \leq r,s \leq n} &= (\sigma(\varphi^\sharp(t_{r,s})))_{1 \leq r,s \leq n} \\ &= \left(\varphi^\sharp \left(\lambda_{\rho(\sigma)}^\sharp(t_{r,s}) \right) \right)_{1 \leq r,s \leq n}, \end{aligned}$$

because φ is equivariant, and

$$\left(\lambda_{\rho(\sigma)}^\sharp(t_{r,s}) \right)_{1 \leq r,s \leq n} = (t_{r,s})_{1 \leq r,s \leq n} a(\rho(\sigma))$$

where for any $\tau \in \mathfrak{V}$ the matrix $a(\tau)$ is in $\mathrm{GL}_n(C)$ and is such that the formula $(\tau(t_{r,s}))_{1 \leq r,s \leq n} = (t_{r,s})_{1 \leq r,s \leq n} a(\tau)$ defines the images of the $t_{r,s}$ by the comorphism $\lambda_\tau^\sharp$ of the right translation on $\mathfrak{V}$ by τ . We have find

$$(\sigma(z_{r,s}))_{1 \leq r,s \leq n} = (z_{r,s})_{1 \leq r,s \leq n} a(\rho(\sigma))$$

with $a(\rho(\sigma))$ in $\mathrm{GL}_n(C)$. $\square$

The fact that φ is an isomorphism implies that $R \otimes_K \tilde{K}$ is generated over $\tilde{K}$ by the $z_{r,s}$, $1 \leq r, s \leq n$, indeed $R \otimes_K \tilde{K}$ is generated over $\tilde{K}$ by the C -space $V := \sum_{1 \leq r,s \leq n} C z_{r,s}$ and the lemma shows that this space V is (globally) invariant under the action of the Galois group $\mathfrak{U}$. The (ordinary) Galois group $\mathrm{Gal}(\tilde{K}/K)$ acts as usual on the right

hand factor of $R \otimes_K \tilde{K}$ and trivially on the left one, then we see that R is generated over K by the $z_{r,s}$, $1 \leq r, s \leq n$.

Another consequence of the previous lemma is that the matrix

$$D \stackrel{\text{def}}{=} (z'_{r,s})_{1 \leq r, s \leq n} (z_{r,s})_{1 \leq r, s \leq n}^{-1},$$

is in $M_n(K)$, then, because $\varphi^\sharp$ is an isomorphism, the ring R is generated by the entries of a fundamental matrix of the equation $Y' = DY$, we know also that R is a simple differential ring. It follows that R , resp. E , is the Picard-Vessiot ring, resp. field, over K of this equation.

To a rational representation $\rho : dGal(L/K) \rightarrow GL_n(C)$ we have associated an element $[D]$ of $Z_n(L/K)$, this is clearly the inverse map of $[A] \mapsto \rho_{[A]}$. $\square$

2. A CORRESPONDANCE.

Let K^{diff} be a universal Picard-Vessiot extension of K and set $G^{\text{diff}} = dGal(K^{\text{diff}}/K)$. We choose once of all an identification $GL_n(C) = GL(C^n)$.

Let $\text{Rep}_n(G^{\text{diff}})$ be the category of representations of G^{diff} in $GL_n(C)$: the objects are morphisms $\rho : G^{\text{diff}} \rightarrow GL_n(C)$, an arrow $f : \rho_1 \rightarrow \rho_2$ is a C -linear map from C^n into itself such that, for any $g \in G^{\text{diff}}$, the following diagram is commutative

$$\begin{array}{ccc} C^n & \xrightarrow{\rho_1(g)} & C^n \\ f \downarrow & & \downarrow f \\ C^n & \xrightarrow{\rho_2(g)} & C^n \end{array}$$

To define the category $\underline{Z}_n(K)$ we need the following remarks. Let M and N be two elements of $M_n(K)$, we say that they are equivalent if there exists U and V in $GL_n(K)$ such that $N = VMU$. We denote by $\overline{M}$ the equivalent class of M . Let $A_i \in M_n(K)$, $i = 1, 2$ and let $M \in M_n(K)$ such that

$$(4) \quad M' = A_2 M - M A_1.$$

Let $B_i \in [A_i]$, let $U_i \in GL_n(K)$ such that

$$A_i = U_i' U_i^{-1} + U_i B_i U_i^{-1},$$

then an easy calculation shows that

$$(U_2^{-1} M U_1)' = B_2 (U_2^{-1} M U_1) - (U_2^{-1} M U_1) B_2.$$

Suppose that $M \in GL_n(K)$ and satisfies (4), then

$$(M^{-1})' = A_1 M^{-1} - M^{-1} A_2.$$

Now we can define the category $\underline{Z}_n(K)$. Its objects are elements of $Z_n(K)$ (see (1)), an arrow $[A_1] \rightarrow [A_2]$, where A_1 and A_2 are elements of $M_n(K)$, is an equivalence class $\overline{M}$ in $M_n(K)$ such that there exists $M \in \overline{M}$ satisfying (4). The two preceding formulas show that this definition does not depend on the choice of A_i in $[A_i]$, $i = 1, 2$, and

that invertible arrows in $\underline{Z}_n(K)$ correspond to equivalence classes of invertible matrices. We explain the composition of arrows. Let $\overline{M} : [A_1] \rightarrow [A_2]$ and $\overline{N} : [A_2] \rightarrow [A_3]$ two arrows of $\underline{Z}_n(K)$, choose $M \in \overline{M}$, $N \in \overline{N}$ such that

$$M' = A_2M - MA_1 \quad \text{and} \quad N' = A_3N - NA_2,$$

then we see that

$$(NM)' = A_3NM - NMA_1.$$

The composed arrow is $\overline{N} \circ \overline{M} = \overline{NM}$, for a good choice of representing elements of the different classes of matrices.

Then $\underline{Z}_n(K)$ is a category, indeed it is easily to see that it is an additive category.

Theorem 2.1. *The two categories $\underline{Z}_n(K)$ and $\underline{\text{Rep}}_n(G^{\text{diff}})$ are equivalent. On objects, this equivalence is $[A] \mapsto c_{[A]}$ (see (1.0.1)).*

Proof. Note that here to write $c_{[A]}$ is an abuse of notation, if L/K is the Picard-Vessiot extension (contained in K^{diff}) associated to the equation $Y' = AY$, we denote always $c_{[A]}$ the representation

$$G^{\text{diff}} \xrightarrow{\text{restriction}} d\text{Gal}(L/K) \xrightarrow{c_{[A]}} \text{GL}_n(C).$$

The map $[A] \mapsto c_{[A]}$ on objects of the categories has been constructed in the previous theorem, it is one to one. Let $[A_1]$ and $[A_2]$ be two objects of $\underline{Z}_n(K)$ and $\overline{M} : [A_1] \rightarrow [A_2]$ be an arrow, select $M \in \overline{M}$ such that $M' = A_2M - MA_1$. Let $F_1, F_2 \in \text{GL}_n(K^{\text{diff}})$ be fundamental matrices for respectively the equations $Y' = A_1Y$ and $Y' = A_2Y$. Then $F'_i = A_iF_i$, $i = 1, 2$. Let $f = F_2^{-1}MF_1$, a priori f is in $\text{GL}_n(K)$, but

$$\begin{aligned} f' &= (F_2^{-1})'MF_1 + F_2^{-1}M'F_1 + F_2^{-1}MF'_1 \\ &= (-F_2^{-1}A_2)MF_1 + F_2^{-1}(A_2M - MA_1)F_1 \\ &\quad + F_2^{-1}MA_1F_1 = 0. \end{aligned}$$

Then $f = F_2^{-1}MF_1$ is in $\text{GL}_n(C)$. Now we prove that f is a morphism from $c_{[A_1]}$ to $c_{[A_2]}$. Let g be an element of G^{diff} . Applying g to the relation $f = F_2^{-1}MF_1$ we find

$$f = g(F_2^{-1})Mg(F_1) = g(F_2^{-1})F_2fF_1^{-1}g(F_1) = c_{[A_2]}(g)^{-1}fc_{[A_1]}(g),$$

(see (1.0.1)) for all g . This means that $f : c_{[A_1]} \rightarrow c_{[A_2]}$ is a map in $\underline{\text{Rep}}_n(G^{\text{diff}})$.

Conversely let $f : \rho_1 \rightarrow \rho_2$ be an arrow of $\underline{\text{Rep}}_n(G^{\text{diff}})$, then we can see f as a matrix with coefficient in C . We know that there exists A_i in $M_n(K)$ such that $\rho_i = c_{[A_i]}$, $i = 1, 2$. Let as before F_i be a fundamental matrix for the equation $Y' = A_iY$. Set $M = F_2fF_1^{-1}$.

- We prove that M is in $M_n(K)$. The fact that f is a morphism of representations means that for all g in G^{diff} we have

$$fc_{[A_1]}(g) = c_{[A_2]}(g)f,$$

which is equivalent to

$$fF_1^{-1}g(F_1) = F_2^{-1}g(F_2)f,$$

then

$$F_2fF_1^{-1} = g(F_2)fg(F_1^{-1}) = g(F_2fF_1^{-1}).$$

This prove that the entries of M are in K .

- We prove the formula $M' = A_2M - MA_1$. We have

$$M' = F_2'fF_1^{-1} + F_2f(F_1^{-1})' = A_2F_2fF_1^{-1} + F_2f(-F_1^{-1}A_1)$$

which is the expected formula. □

REFERENCES

- [1] van der PUT Marius - REVERSAT Marc. Arithmetic problems for differential equations (forthcoming).
- [2] van der PUT Marius - SINGER Michael F. Galois theory of linear differential equations. Grundlehren der Mathematischen Wissenschaften 328, Springer Verlag 2003.

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UNIVERSITÉ PAUL SABATIER,
118 ROUTE DE NARBONNE, 31062 TOULOUSE CEDEX 9, FRANCE.

E-mail address: marc.reversat@math.ups-tlse.fr